

Holoprosencephaly: New Concepts

Erin M Simon, MD, OTR, Assistant Professor of Radiology, The Children's Hospital of Philadelphia, Department of Radiology, 34th Street and Civic Center Boulevard, Philadelphia, Pennsylvania 19104-4399, Office: 215.590.1030 Fax: 215.590.1345

Since the introduction of the term “arhinencephaly” by Hans Kundrat (1882), investigators have attempted to characterize and further understand the broad spectrum of malformations resulting from a lack of separation of the structures of the forebrain. These malformations are generally lumped under the term “holoprosencephaly” (HPE) coined by DeMyer and Zeman in 1963. These authors further subdivided HPE into the classifications of alobar, semilobar and lobar. More recent work has described a form of HPE in which the basal forebrain is relatively spared and the posterior portion of the frontal lobe is non-cleaved, the middle interhemispheric variant.

While there has been a vast amount of attention in the literature devoted to the complex facial malformations that often accompany the brain parenchymal abnormalities in HPE, it is the generally profound brain anomalies that dominate the clinical picture and prognosis. While the “face predicts the brain” in approximately 80% of cases, it is the brain that predicts the outcome in all. Expanding our knowledge of the extensive brain malformations will allow us to better serve these patients and their families.

Recent discoveries in the fields of genetics and developmental neurobiology have also advanced our knowledge of HPE. The first descriptions of mutations in the gene Sonic hedgehog (Shh) as a causative agent in HPE have given way to the identification and implication of many other gene mutations in the formation of this complex spectrum of disorders. Great strides have been made in understanding human central nervous system development from extrapolating information obtained in cellular fate maps of several animal species. By combining this basic science knowledge with our observations of the brain morphology from imaging studies of a large patient cohort, we can begin to better understand the factors that contribute to the formation of this fascinating spectrum of forebrain malformation. We present evidence that an imbalance of dorsal and ventral patterning signals in the secondary prosencephalon and floor plate result in many of the morphological abnormalities seen in the more common forms of HPE. Ultimately, we hope to further correlate the morphology and biology with neurodevelopmental information, in order to establish more accurate classification of the spectrum and stratification measures for prediction of patient outcome.

Simon EM, Barkovich AJ. Holoprosencephaly: new concepts. *Magnetic Resonance Imaging Clinics of North America; Pediatric MR Neuroimaging*. W.B.Saunders Company, Philadelphia, PA, 2001;9:1,149-164

References

Simon EM, Hevner RF, Pinter JD, Clegg NJ, Miller VS, Kinsman SL, Hahn JS, Barkovich AJ. Assessment of the deep gray nuclei in holoprosencephaly, *AJNR*, *Am J Neuroradiol* 2000;21:1955-1961 (PDF can be downloaded at no charge via www.ajnr.org)

Barkovich AJ, **Simon EM**, Walsh CA. Agenesis of the corpus callosum with interhemispheric cyst: review of 25 cases and structural analyses, *Neurology* 2001;56:220-227

Simon EM, Hevner RF, Pinter JD, Clegg NJ, Delgado M, Kinsman SL, Hahn JS, Barkovich AJ. The dorsal cyst in holoprosencephaly and the role of the thalami in its formation, *Neuroradiology* 2001;9:787-791

Simon EM, Hevner RF, Pinter JD, Clegg NJ, Delgado M, Kinsman SL, Hahn JS, Barkovich AJ. The middle interhemispheric variant of holoprosencephaly. *AJNR Am J Neuroradiol*, 2002;23:151-155 (PDF can be downloaded at no charge via www.ajnr.org)

Barkovich AJ, **Simon EM**, Clegg NJ, Kinsman SL, Hahn JS. Analysis of the cerebral cortex in holoprosencephaly with attention to the sylvian fissures. *AJNR Am J Neuroradiol*, 2002;23:143-150 (PDF can be downloaded at no charge via www.ajnr.org)